

Automorphisms of a family of surfaces with $p_g = q = 2$ and $K^2 = 7$

Matteo Penegini and Roberto Pignatelli

In memoria di Gianfranco, un amico e un maestro

Abstract. We compute the automorphism group of all the elements of a family of surfaces of general type with $p_g = q = 2$ and $K^2 = 7$, originally constructed by C. Rito in [Rit18]. We discuss the consequences of our results towards the Mumford-Tate conjecture.

1. Introduction

The classification of general type surfaces with low numerical invariants is unanimously considered a very difficult problem to tackle. It is already difficult to construct new surfaces with low numerical invariants. Therefore, as soon as new examples are found, it is natural to test the famous conjectures on them. This short note stems from the question whether it is possible to verify the Mumford-Tate conjecture for surfaces S of general type with $p_g = q = 2$ and $K^2 = 7$ first constructed by Rito in [Rit18] and later studied by the authors in [PePi20].

These surfaces S are obtained as a generically finite double covering of an abelian surface A , which turns out to be the Albanese variety of S , branched along a curve with a singular point of type $(3, 3)$ and no other singularities. These surfaces give rise to three disjoint open subsets in the Gieseker moduli space $\mathcal{M}_{2,2,7}^{\text{can}}$ which are all irreducible, generically smooth of dimension 2, that we shall denote by \mathcal{M}_1 , \mathcal{M}_2 and \mathcal{M}_4 .

The Mumford-Tate conjecture for surfaces is still an open problem and only in very few examples has it been verified as true, see for example [Moo17a]. A strategy to prove the conjecture for surfaces with $p_g = q = 2$ and of maximal Albanese dimension is outlined in the article [CoPe20]. The strategy reduces to finding geometric quotients X of S that are K3 surfaces whose weight 2 Hodge structure is a sub-Hodge structure of the weight 2 Hodge structure of S orthogonal to the the sub-Hodge structure coming from the Albanese surface of S . This strategy proved to be very successful in many cases, see [CoPe20]. The first question toward exploiting the strategy is to calculate the automorphism of the surfaces S and then classify all possible quotients. This is the content of the main theorem of this note.

THEOREM 1.1. The automorphism group of the surface S of general type with $p_g = q = 2$ and $K^2 = 7$ constructed by Rito in [Rit18] is a product of cyclic groups

$$\text{Aut}(S) \cong G \times \mathbb{Z}/2\mathbb{Z} = G \times \langle \sigma \rangle,$$

where $S/\langle \sigma \rangle \cong A = \text{Alb}(S)$, while

1. G is trivial if $[S] \in \mathcal{M}_1$;
2. $G \cong \mathbb{Z}/2$ if $[S] \in \mathcal{M}_2$;
3. $G \cong \mathbb{Z}/4$ if $[S] \in \mathcal{M}_4$.

The second step of the strategy is to identify the quotients. We have at once

COROLLARY 1.2. For all $H \leq \text{Aut}(S)$, the quotients $X = S/H$ are irregular surfaces, i.e., $q(X) \geq 1$.

This corollary tells us that in order to prove the Mumford Tate conjecture for these surfaces a new strategy is needed.

Now, let us explain the way in which this paper is organized.

In the second section we recall the construction of the surfaces S with the calculation of the invariants. The third section is devoted to the proof of the Main Theorem. The section four contains the calculation of the invariants of the quotient surfaces. Finally we include a section five where it is explained the strategy to prove the conjecture for surfaces with $p_g = q = 2$ and of maximal Albanese dimension.

2020 Mathematics Subject Classification: 14J29, 14J10, 14B12

Keywords: Surface of general type, Albanese map, Automorphisms of varieties

Version: March 4, 2024

Acknowledgments. Both authors were partially supported by GNSAGA-INdAM. The first author was partially supported by GNSAGA-INdAM and PRIN 2020KKWT53 003 - Progetto: *Curves, Ricci flat Varieties and their Interactions* and partially supported by the DIMA - Dipartimento di Eccellenza 2023-2027.

Notation and conventions. We work over the field \mathbb{C} of complex numbers. By *surface* we mean a projective, non-singular surface S , and for such a surface K_S denotes the canonical class, $p_g(S) = h^0(S, K_S)$ is the *geometric genus*, $q(S) = h^1(S, K_S)$ is the *irregularity* and $\chi(O_S) = 1 - q(S) + p_g(S)$ is the *Euler-Poincaré characteristic*.

2. The surfaces

In this section we report, for the convenience of the reader, the construction of the families of surfaces under consideration. We use the notation of [PePi20], the main result in this direction is the following one.

PROPOSITION 2.1. [PePo13b] Let A be an Abelian surface. Assume that A contains a reduced curve whose class is 2-divisible in $\text{Pic}(A)$, whose self intersection is 16, with a unique singular point of type $(3,3)$ and no other singularity. Then there exists a generically finite double cover $S \rightarrow A$ branched along this curve. Moreover, the numerical invariants of S are $p_g(S) = q(S) = 2$ and $K_S^2 = 7$.

In [Rit18] and later in [PePi20] the existence of the abelian surface A , that has the properties as in Proposition 2.1, is proved. In particular, it is also shown that the double cover coincides with the Albanese map, hence A is the Albanese variety associated to S , we denote it by

$$\alpha: S \rightarrow \text{Alb}(S) = A.$$

We can be more precise, A is isogenous to a product of two elliptic curves T_1 and T_2 . We denote by

$$\iota: A \rightarrow T_1 \times T_2$$

the isogeny, which is of degree 2. Clearly, A carries a $(1,2)$ -polarization L which is a pull-back of a (product) principal polarization via the isogeny ι . In addition, on A we have two elliptic fibrations $f_j: A \rightarrow T_j$ with fibres $\Lambda_j: \Lambda_1 \cong T_2$ and $\Lambda_2 \cong T_1$.

The branching locus of α is an effective divisor with two irreducible components

$$C_1 + t \in |2L|,$$

where $C_1 = f_2^{-1}(b_1)$ is an element of $|\Lambda_2|$, with $b_1 \in T_1$. While, t is a curve of geometric genus 3 with a tacnode tangent to C_1 at a point p . The situation is exemplified in the following Figure 1.

We deduce that

$$(1) \quad C_1^2 = 0, \quad t^2 = 8, \quad C_1 t = 4,$$

notice that $(C_1 + t)^2 = 16$.

Notice that the branch locus is singular in p . Therefore, to get a smooth surface S as a generically finite double cover of A branched along $C_1 + t$ we have to blow up the point p (see Figure 1) first.

1. First, we resolve the singularity in p . To do that, we need to blow up A twice, first in p and then in a point infinitely near to p . Let us denote these two blow ups by

$$B' \xrightarrow{\sigma_4} B \xrightarrow{\sigma_3} A.$$

On B' , let us denote by F the exceptional divisor relative to σ_4 , by E' the strict transform of the exceptional divisor E relative to σ_3 , by C_1 the strict transform of C_1 and, finally, by R the strict transform of t (see Figure 1).

In addition, one gathers the following information: $E' \cong \mathbb{P}^1$ and $(E')^2 = -2$, $F \cong \mathbb{P}^1$ and $F^2 = -1$, $g(C_1) = 1$ and $C_1^2 = -2$.





Figure 1: The Branch Divisor of α and β

2. Second, we consider a double cover of $\beta: S' \rightarrow B'$ ramified over $R + C_1 + E'$ (this is even since $t + C_1$ is even on A). The surface S' is a surface of general type, not minimal. Indeed, it contains a -1 -curve, which is $\hat{E} = \beta^{-1}(E')$. The ramification divisor is denoted $\hat{R} + \hat{C}_1 + \hat{E}$. Notice that \hat{C}_1 has genus 1 and $\hat{C}_1^2 = -1$.
3. Finally, to get S we contract the -1 -curve \hat{E} .

We can summarize the construction of S with the following diagram.



Moreover, the point p is a $[3,3]$ point, which is not a negligible singularity. A $[3,3]$ point is a pair (x_1, x_2) such that x_1 belongs to the first infinitesimal neighborhood of x_2 and both are triple points for the curve. Thus, we may calculate the invariants of S by using the formulae in [BHPV03, p. 237]. In those formulae x_2 counts as a triple point (so $m_2 = 1$) and x_1 as a quadruple point (so $m_1 = 2$). Then

$$(2) \quad 2 = 2\chi(O_{S'}) = L^2 - \sum_{i=1}^2 m_i(m_i - 1), \quad K_{S'}^2 = 2L^2 - 2 \sum_{i=1}^2 (m_i - 1)^2,$$

which imply

$$K_S^2 \geq K_{S'}^2 = 4 + 2 \sum_{i=1}^2 (m_i - 1).$$

Finally, once we contract the -1 -curve on S' , we obtain hence $K_S^2 = 7$ and $\chi(S) = 1$.

Considering the Abelian varieties $A, T_1, T_2, T_1 \times T_2$ we choose the following points as neutral elements:

$$p \in A, \quad a_3 := f_1(p) \in T_1, \quad b_1 := f_2(p) \in T_2, \quad (a_3, b_1) \in T_1 \times T_2.$$

With this particular choice ι, f_1, f_2 are homomorphism of groups too.

The remaining 2-torsion points on each elliptic curve T_j will be denoted by

$$a_1, a_2, a_4 \in T_1[2], \quad b_2, b_3, b_4 \in T_2[2].$$

This yields $\iota_* O_A^- \cong O_{T_1}(a_4 - a_3) \boxtimes O_{T_2}(b_2 - b_1)$, where $\iota_* O_A^-$ is the anti-invariant part of $\iota_* O_A$, see [PePi20, Lemma 3.4] for a detailed proof.

Remark 2.2. Furthermore in [PePi20] it is proved that

$$f_1^*(a_4 + a_3) + f_2^*(b_3 + b_1) \in |2L|,$$

whence

$$L \cong f_1^*(\bar{a}) \otimes f_2^*(\bar{b})$$

where \bar{b} is a 4-torsion point such that $\bar{b} \oplus \bar{b} \neq b_2$. While for \bar{a} we have three possible choices

1. $\bar{a} = a_3$ (in this case $\bar{b} \oplus \bar{b} = b_4$);
2. \bar{a} is a 2-torsion point such that $\bar{a} \neq a_4$ (in this case $\bar{b} \oplus \bar{b} = b_4$);
3. \bar{a} is a 4-torsion point such that $\bar{a} \oplus \bar{a} \neq a_4$ (in this case $\bar{b} \oplus \bar{b} = b_3$).

As just remarked all the choices are possible and to each choice corresponds a different irreducible component of the Gieseker moduli space $\mathcal{M}_{2,2,7}^{\text{can}}$ of the canonical models of the surfaces of general type with $p_g = q = 2$ and $K^2 = 7$. We shall denote these components, following [PePi20, Proposition 5.8], by $\mathcal{M}_i \subset \mathcal{M}_{2,2,7}^{\text{can}}$ with $i \in \{1, 2, 4\}$ respectively. Note that the index i equals the order of \bar{a} as torsion point.

3. The automorphisms of the Rito's surfaces

Consider the abelian variety A . We know that there is an isogeny of degree 2 onto a product of elliptic curves $T_1 \times T_2$.

By taking the universal covers, we can write $T_j := \mathbb{C}/\lambda_j$ where the $\lambda_j \cong \mathbb{Z}^2$ are lattices so that the origin maps to a_3 respectively b_1 . We choose generators \bar{e}_1, \bar{e}_2 of λ_1 , \bar{e}_3, \bar{e}_4 of λ_2 so that $\frac{\bar{e}_1}{2}$ maps to a_1 , $\frac{\bar{e}_2}{2}$ maps to a_2 , $\frac{\bar{e}_3}{2}$ maps to b_3 , $\frac{\bar{e}_4}{2}$ maps to b_4 .

So, in \mathbb{C}^2 -coordinates we have

$$e_1 = (\bar{e}_1, 0), \quad e_2 = (\bar{e}_2, 0), \quad e_3 = (0, \bar{e}_3), \quad e_4 = (0, \bar{e}_4).$$

Since the universal cover of $T_1 \times T_2$ factors through the isogeny ι , we obtain $A = \mathbb{C}^2/\lambda$ where λ is a sublattice of index 2 of the lattice

$$\lambda_1 \oplus \lambda_2 = \left\{ \sum t_i e_i \mid t_i \in \mathbb{Z} \right\}$$

LEMMA 3.1. The lattice λ is the sublattice of $\lambda_1 \oplus \lambda_2$ of the elements whose sum $\sum t_i$ is even.

Proof. We set $\tilde{\lambda} := \left\{ \sum m_j e_j \mid \sum m_j \text{ is even} \right\}$, $\tilde{A} := \mathbb{C}^2/\tilde{\lambda}$. The inclusion $\tilde{\lambda} \subset \lambda_1 \oplus \lambda_2$ induces an isogeny $\tilde{\iota}: \tilde{A} \rightarrow T_1 \times T_2$ of degree 2.

An isogeny of degree 2 is determined by the anti-invariant part of the direct image of the trivial bundle, that is a generator of the kernel of the pull-back map among the Picard groups. So we only need to prove that $\tilde{\tau}^*(\mathcal{O}_{T_1}(a_4 - a_3) \boxtimes \mathcal{O}_{T_2}(b_2 - b_1))$ is the trivial sheaf of \tilde{A} .

It is then enough to show that the line bundles $\mathcal{O}_{T_1}(a_4 - a_3)$ and $\mathcal{O}_{T_2}(b_2 - b_1)$ pull back to the trivial bundle of \tilde{A} via the natural fibrations $\tilde{A} \rightarrow T_j$. We prove it only for $\mathcal{O}_{T_2}(b_2 - b_1)$, the other case being perfectly analogous.

Let $\lambda'_2 \subset \lambda_2$ be the index 2 sublattice of the elements of the form $m_3\bar{e}_3 + m_4\bar{e}_4$ with $m_3 + m_4$ even. This defines an isogeny of degree 2, $\tilde{\iota}_2: \mathbb{C}/\lambda'_2 \rightarrow T_2$, fitting in the commutative diagram below.

$$(3) \quad \begin{array}{ccc} \tilde{A} & \xrightarrow{\tilde{\iota}} & T_1 \times T_2 \\ \downarrow f'_2 & & \downarrow \\ \mathbb{C}/\lambda'_2 & \xrightarrow{\tilde{\iota}_2} & T_2 \end{array}$$

It is then enough to show the triviality of $\tilde{\iota}_2^*\mathcal{O}_{T_2}(b_2 - b_1)$ on \mathbb{C}/λ'_2 . This is standard; it can be show for example as follows.

The point b_1 pulls back to the sum of two points, the classes modulo λ'_2 of 0 and \bar{e}_3 . The point b_2 pulls back to the sum of the classes of $\frac{1}{2}(\bar{e}_3 + \bar{e}_4)$ and $\bar{e}_3 + \frac{1}{2}(\bar{e}_3 + \bar{e}_4)$. Since

$$\left(\frac{1}{2}(\bar{e}_3 + \bar{e}_4) + \bar{e}_3 + \frac{1}{2}(\bar{e}_3 + \bar{e}_4) \right) - (0 + \bar{e}_3) = \bar{e}_3 + \bar{e}_4 \in \lambda'_2$$

the two divisors of degree 2 we have obtained are linearly equivalent. \square

We will need the following general result for an abelian surface with a $(1, 2)$ -polarization, the proof of which can be found in [Bar87, Section 1.2].

Remark 3.2. The linear system $|L|$ contains exactly two reducible divisors union of elements respectively of Λ_1 and Λ_2 , the curves $f_1^*\bar{a} + f_2^*\bar{b}$ and $f_1^*(\bar{a} \oplus a_4) + f_2^*(\bar{b} \oplus b_2)$.

Since the Albanese morphism $\alpha: S \rightarrow A$ has degree 2, it determines an involution $\sigma: S \rightarrow S$ that is central in $\text{Aut } S$ and an exact sequence

$$0 \rightarrow \mathbb{Z}/2\mathbb{Z} = \langle \sigma \rangle \rightarrow \text{Aut } S \rightarrow G \rightarrow 0$$

where G is the group of the self-biholomorphisms $\varphi: A \rightarrow A$ such that

1. $\varphi^*L = L$;
2. $\varphi(C_1 + t) = C_1 + t$, equivalently $\varphi(C_1) = C_1$, $\varphi(t) = t$, $\varphi(p) = p$.

We have written an isomorphism $A \cong \mathbb{C}^2/\lambda$ where the point p is the image of the origin of \mathbb{C}^2 . From $\varphi(p) = p$ it follows that the elements of G are automorphisms of A as a group with the group structure induced by \mathbb{C}^2 . Now, we can see that one of the above assumption is not necessary, indeed we have the following lemma.

LEMMA 3.3. The group G is the group of the automorphisms of the Abelian variety A preserving the group structure induced by the identification $\mathbb{C}^2/\lambda = A$ such that $\varphi^*L = L$ and $\varphi(C_1) = C_1$.

Proof. The only nontrivial thing to prove is that, if $\varphi: A \rightarrow A$ is a group automorphism such that $\varphi^*L = L$ and $\varphi(C_1) = C_1$, then $\varphi(t) = t$.

By (1), the intersection number between t and C_1 is 4. More precisely, t cuts on C_1 the divisor $4p$. Since $\varphi(p) = p$ and $\varphi(C_1) = C_1$, $\varphi(t)$ cuts $4p$ on C_1 as well.

Assume by contradiction $\varphi(t) \neq t$. Then the pencil generated by $\varphi(t)$ and t contains a divisor of the form $C_1 + D_1$, where D_1 is an effective divisor in $|t - C_1| = |2L - 2C_1|$.

Let D be any element of Λ_2 .

The map f'_2 , in the commutative diagram (3), maps D isomorphically to \mathbb{C}/λ'_2 and the divisor cut by D_1 on D to a divisor in the linear system $|\tilde{\iota}_2^*(2\bar{b} - 2b_1)|$. The line bundle $\mathcal{O}_{T_2}(2\bar{b} - 2b_1)$ is of 2-torsion. Since $\bar{b} \oplus \bar{b} \neq b_2$, the

proof of Lemma 3.1 shows that $\tilde{\tau}_2^* \mathcal{O}_{T_2}(2\bar{b} - 2b_1)$ is not trivial and then $|\tilde{\tau}_2^*(2\bar{b} - 2b_1)|$ is empty. Then D_1 contains D . Varying D in Λ_2 we obtain $D_1 = \bar{0}$, a contradiction. \square

The action of G on A may be uniquely lifted to an action on \mathbb{C}^2 fixing the origin, so representing G as a finite subgroup of the linear group $GL_2(\mathbb{C})$ of the matrices preserving the lattice λ . We will then write ϕ as a matrix

$$\phi = \begin{pmatrix} \phi_{11} & \phi_{12} \\ \phi_{21} & \phi_{22} \end{pmatrix}$$

LEMMA 3.4. G is a group of diagonal matrices. In other words $\phi_{12} = \phi_{21} = 0$.

Proof. We consider \mathbb{C}^2 with the natural coordinates (x, y) given by the construction, so that the line $x = 0$ is the connected component through the origin of the preimage of C_1 in \mathbb{C}^2 . Since $\phi(C_1) = C_1$ the matrix ϕ preserves $x = 0$, so $\phi_{12} = 0$.

Now we use that $\phi^*L = L$. Since by Remark 3.2, $|L|$ contains exactly two reducible divisors, ϕ either preserves or exchange them. The preimage of both divisors on \mathbb{C}^2 is a union of countably many "horizontal" lines $y = c$ and countably many vertical lines $x = c$. However, if $\phi_{21} \neq 0$ any horizontal line is mapped to a line that is neither horizontal nor vertical, a contradiction. \square

In particular ϕ is given by two roots of the unity $\phi_{jj} \in \mathbb{C}$ giving automorphisms of the two elliptic curves: each ϕ_{jj} gives an automorphism of T_j .

We will now need the following well known facts on automorphisms of elliptic curves see for instance [Sil08, Section III.10].

LEMMA 3.5. Let ω be a nontrivial automorphism of an elliptic curve of order n . Then $n = 2, 3, 4$ or 6 . Moreover

1. for every 4-torsion point $p \in T$, $\omega(p) \neq p$;
2. if $n = 3, 6$, then for every 2-torsion point $p \in T$, $\omega(p) \neq p$;
3. if $n = 4$ then there is exactly one 2-torsion point $p \in T$ such that $\omega(p) = p$;
4. if $n = 2$ then $\omega(p) = p$ for all 2-torsion points $p \in T$.

With these facts in mind we can prove

PROPOSITION 3.6. We have the following possibilities for the group G according to the cases in Remark 2.2.

- » In case 1: if T_1 has an automorphism of order 4, G is cyclic of order 4 generated by the automorphism given by $\phi_{11} = i$, $\phi_{22} = 1$, that is

$$(x, y) \mapsto (ix, y).$$

- » In case 3: G is the trivial group of order 1.
- » In the remaining cases, G is a cyclic group of order 2 generated by the automorphism given by $\phi_{11} = -1$, $\phi_{22} = 1$, the involution

$$(x, y) \mapsto (-x, y)$$

Proof. By Lemma 3.4 ϕ acts on the fibrations $f_j: A \rightarrow T_j$ acting on the codomain by ϕ_{jj} .

Since by Remark 3.2, $|L|$ contains exactly two reducible divisors, $f_1^* \bar{a} + f_2^* \bar{b}$ and $f_1^*(\bar{a} \oplus a_4) + f_2^*(\bar{b} \oplus b_2)$, ϕ either preserves or exchange them. So, on $T_1 \times T_2$, the matrix ϕ maps (\bar{a}, \bar{b}) either to (\bar{a}, \bar{b}) or to $(\bar{a} \oplus a_4, \bar{b} \oplus b_2)$.

We now show $\phi_{22} = 1$. In fact in both cases $\phi_{22}^2(\bar{b}) = \bar{b}$. Since \bar{b} is a 4-torsion point, by Lemma 3.5, part 1, $\phi_{22}^2 = 1$. Moreover, if $\phi_{22} \neq 1$ (so $\phi_{22} = -1$) $\phi_{22}(\bar{b}) = \bar{b} \oplus b_2$ that implies $\bar{b} \oplus \bar{b} = b_2$, a contradiction. So $\phi_{22} = 1$.

As a first consequence, $\phi_{11}(\bar{a}) = \bar{a}$.

Now recall that the matrix ϕ preserves the lattice λ . Since $\phi_{22} = 1$, then ϕ_{11} preserves λ_1 and the index 2 sublattice $\lambda'_1 \subset \lambda_1$ of the elements of the form $m_1 \bar{e}_1 + m_2 \bar{e}_2$ with $m_1 + m_2$ even. This implies $\phi_{11}(\bar{e}_1 + \bar{e}_2) - (\bar{e}_1 + \bar{e}_2) \in 2\lambda_1$. Dividing by 2 we obtain $\phi_{11}(a_4) = a_4$.

Now we distinguish the three cases according to Remark 2.2.

In case 3, \bar{a} is a 4-torsion point. Then by $\varphi_{11}(\bar{a}) = \bar{a}$ and Lemma 3.5, part 1, $\varphi_{11} = 1$.

In case 2, \bar{a} and a_4 are distinct 2-torsion points fixed by φ_{11} . Then by Lemma 3.5, part 2 and 3, φ_{11} has order 1 or 2, so $\varphi_{11} = \pm 1$. On the other hand the map $(x, y) \mapsto (-x, y)$ preserves λ so it defines an automorphism of A that defines an element of G .

Finally, case 1. In this case $\varphi_{11}(\bar{a}) = \bar{a}$ holds independently by the choice of the complex number of φ_{11} . Still we have the condition $\varphi_{11}(a_4) = a_4$ that by Lemma 3.5, part 2, forces $\varphi_{11}^4 = 1$. If T_1 has no automorphisms of order 4, then we obtain $\varphi_{11} = \pm 1$ and we conclude as in case 2. Else analogous argument shows that $(x, y) \mapsto (ix, y)$ generates G . \square

Then we can compute $\text{Aut}(S)$.

PROPOSITION 3.7.

$$\text{Aut}(S) \cong G \times \mathbb{Z}/2\mathbb{Z}$$

Proof. Choose any element $g \in G$. Then by 3.6 g acts as $(x, y) \mapsto (kx, y)$ in the coordinates considered there, where k is a complex number with $k^4 = 1$. In those coordinates C_1 is defined by y .

Let Z be the finite double cover of A birational to S , and let $q \in Z$ be the unique point over $p = (0, 0)$. In a neighbourhood of q , Z has equation $z^2 = f(x, y)$ where f is an equation of the branch locus, that is geometrically g -invariant. Then there is a constant c such that $g^*f = cf$. From $f(x, y) = y^3 + O(4)$ we deduce $c = 1$ and $g^*f = f$.

The involution on Z induced by the Albanese morphism of S acts near q as $(x, y, z) \rightarrow (x, y, -z)$; g acts $(x, y) \rightarrow (kx, y)$, so its liftings of g act as $(x, y, z) \mapsto (kx, y, \pm z)$. So the liftings acting locally trivially on the variable z form a splitting map $G \rightarrow \text{Aut}(S)$ mapping to a subgroup that commutes with the Albanese involution. \square

4. The quotients of S

In this brief section we shall prove Corollary 1.2. By Proposition 3.7 $\text{Aut}(S) = \langle \sigma \rangle \times G$ let us consider $H \leq \text{Aut}(S)$, then we have the following diagram

$$\begin{array}{ccc} & S' & \\ & \swarrow & \searrow \\ A & & X \cong S/H \end{array}$$

A natural question to address is the classification of quotient surfaces S/H .

A first step in studying the quotient surfaces is to determine their numerical invariants. To this end we study the induced action of the group H on the cohomology groups of S . Recall that the the global sections of $H^{1,0}(S)$ comes from the one forms on A that we denote by dx, dy . Moreover, one of the two generators of the global sections of $H^{2,0}(S)$ can be identified with $dx \wedge dy$, and of course being $p_g(S) = 2$ we have a global 2-forms ω not coming from A . To summarize this we can write

$$H^{2,0}(S) \cong \langle dx \wedge dy, \omega \rangle, \quad H^{1,0}(S) \cong \langle dx, dy \rangle.$$

Let us denote by g the generator of the cyclic group H , which has order 1, 2 or 4 according to the three cases of Proposition 3.6. The same proposition describes completely the induced action on the generators of the cohomology groups in each case. In particular, we have

$$dx \mapsto g^*dx = \begin{cases} dx \\ -dx \\ idx \end{cases}, \quad dy \mapsto g^*dy = dy.$$

This yields at once that $q(X) \geq 1$, and thus we have proved Corollary 1.2. We can remark already that no quotient X can be a K3 surface. Moreover, we have

$$dx \wedge dy \mapsto \begin{cases} dx \wedge dy \\ -dx \wedge dy, \\ idx \wedge dy. \end{cases}$$

We have two cases for the other generator either $g^*\omega \neq \omega$ or $g^*\omega = \omega$. In the first case the surface X has $p_g(X) = 0$ and in the other cases $p_g(X) = 1$.

All in all by the theorem of classification of algebraic surfaces we have that the quotient X can be either a rational elliptic surface or a properly elliptic surface.

5. Towards the Mumford-Tate conjecture

This section is devoted to explain the strategy used up to know for proving the Mumford Tate conjecture for surfaces with $p_g(S) = q(S) = 2$ and of Albanese maximal dimension.

Let S be a smooth projective complex surface with invariants $p_g(S) = q(S) = 2$, and assume that the Albanese morphism $\alpha: S \rightarrow A$ is surjective. We can make the following general observations (see also [CoPe20]). It holds:

1. The induced map on cohomology $\alpha^*: H^*(A, \mathbb{Z}) \rightarrow H^*(S, \mathbb{Z})$ is injective. The orthogonal complement $H_{\text{new}}^2 = H^*(A, \mathbb{Z})^\perp \subset H^*(S, \mathbb{Z})$ is a Hodge structure of weight 2 with Hodge numbers $(1, n, 1)$, where $n = h^{1,1}(S) - 4$. Such a Hodge structure is said to be of *K3 type*.
2. Let S' be a smooth projective complex surface with invariant $p_g(S') = 1$. Then Morrison [Mor87] showed that there exists a K3 surface X' together with an isomorphism $\iota': H^2(S', \mathbb{Q})^{\text{tra}} \rightarrow H^2(X', \mathbb{Q})^{\text{tra}}$ that preserves the Hodge structure, the integral structure, and the intersection pairing. (Here $(_)^{\text{tra}}$ denotes the *transcendental* part of the Hodge structure, that is, the orthogonal complement of the Hodge classes.)

We now look closely to our surfaces S with $p_g(S) = q(S) = 2$, for which we know that the Albanese map is a generically finite cover.

Then we have

PROPOSITION 5.1. Let S be a smooth projective complex surface with invariants $p_g(S) = q(S) = 2$, and assume that the Albanese morphism $\alpha: S \rightarrow A$ is surjective. Then there exists a K3 surface X and an isomorphism of Hodge structures

$$\iota: (H_{\text{new}}^2(S, \mathbb{Q}))^{\text{tra}} \rightarrow H^2(X, \mathbb{Q})^{\text{tra}}.$$

This is a direct consequence of [Mor87]. Notice that the surface X is related only Hodge theoretically to S . Therefore, this is not enough to prove the conjecture, to this end we have to address the following question:

Do there exist X and ι as above, such that ι is motivated in the sense of Andrè?

Let us briefly explain and recall some facts on categories of motives, and for the reader convenience we state the motivic Mumford–Tate conjecture, for a more detailed introduction on the subject see [Moo17b, Moo17a]. First we recall some facts about Chow motives and André motives of surfaces. We do not need full generality, so let K be a subfield of \mathbb{C} .

Given smooth and projective varieties X and Y over a field K (i.e., objects in the category SmPr_K) of dimension d_X and d_Y respectively, a correspondence of degree k from X to Y is an element γ of $A^{d_X+k}(X \times Y)$. Then γ induces a map $A^*(X) \rightarrow A^{*+k}(Y)$ by the formula

$$\gamma_*(\beta) := \pi_{2*}(\gamma \cdot \pi_1^*(\beta)),$$

where $\pi_1: X \times Y \rightarrow X$ and $\pi_2: X \times Y \rightarrow Y$ denote the projections. The category \mathcal{M}_{rat} of *Chow motives* (with rational coefficients) over K is defined as follows:

- » the objects of \mathcal{M}_{rat} are triples (X, p, n) such that $X \in \text{SmPr}_{/\mathbb{C}}$, $p \in A^{d_X}(X \times X)$ is an idempotent correspondence (i.e. $p_* \circ p_* = p_*$) and n is an integer;

» the morphisms in \mathcal{M}_{rat} from (X, p, n) to (Y, q, m) are correspondences $f: X \rightarrow Y$ of degree $n - m$, such that $f \circ p = f = q \circ f$.

We recall that \mathcal{M}_{rat} is an additive, \mathbb{Q} -linear, pseudoabelian category, see [?, Theorem 1.6].

There exists a functor

$$h: \text{SmPr}_{/\mathbb{C}}^{\text{op}} \rightarrow \mathcal{M}_{\text{rat}} \text{ such that } h: X \mapsto (X, 1, 0)$$

from the category of smooth projective varieties over \mathbb{C} to the category of Chow motives.

We denote also with $A^i(M)$ the i -th Chow group of a motive $M \in \mathcal{M}_{\text{rat}}$. In general, it is not known whether the Künneth projectors π_i are algebraic, so it does not (yet) make sense to speak of the summand $h^i(X) \subset h(X)$ for an arbitrary smooth projective variety X/\mathbb{C} . However, a so-called Chow–Künneth decomposition does exist for curves [Man68], for surfaces [Mur90], and for abelian varieties [DeMu91]. For algebraic surfaces there is in fact the following theorem, which strengthens the decomposition of the Chow motive. Statement and proof are copied from [Lat19, Theorem 2.2].

THEOREM 5.2. Let S be a smooth projective surface over \mathbb{C} . There exists a self-dual Chow–Künneth decomposition $\{\pi_i\}$ of S , with the further property that there is a splitting

$$\pi_2 = \pi_2^{\text{alg}} + \pi_2^{\text{tra}} \in A^2(S \times S)$$

in orthogonal idempotents, defining a splitting $h^2(S) = h_{\text{alg}}^2(S) \oplus h_{\text{tra}}^2(S)$ with Chow groups

$$A^i(h_{\text{alg}}^2(S)) = \begin{cases} \text{NS}(S) & \text{if } i = 1, \\ 0 & \text{otherwise,} \end{cases} \quad \text{and} \quad A^i(h_{\text{tra}}^2(S)) = \begin{cases} A_{\text{AJ}}^2(S) & \text{if } i = 2, \\ 0 & \text{otherwise.} \end{cases}$$

Here $A_{\text{AJ}}^2(S)$ denotes the kernel of the Abel–Jacobi map.

The idea to construct such K3 surface X is to exploit the automorphism group G of S and prove that a quotient S/H by some subgroup $H \leq G$ is birational to X . Of course this will give us a weak answer to the previous question. Nevertheless, it will suffice to prove that – using the notion of *motivated cycles* introduced by André [And96] – there exist X and \mathfrak{t} as above, such that \mathfrak{t} is motivated.

To speak about motivic Mumford–Tate conjecture we need to introduce the notion of *motivated cycles* (for a brief introduction see e.g. [Moo17b, Section 3.1]).

Definition 5.3. Let K be a subfield of \mathbb{C} , and let X be a smooth projective variety over K . A class γ in $H^{2i}(X)$ is called a *motivated cycle* of degree i if there exists an auxiliary smooth projective variety Y over K such that γ is of the form $\pi_*(\alpha \cup \star\beta)$, where $\pi: X \times Y \rightarrow X$ is the projection, α and β are algebraic cycle classes in $H^*(X \times Y)$, and $\star\beta$ is the image of β under the Hodge star operation.

Every algebraic cycle is motivated, and under the Lefschetz standard conjecture the converse holds as well. The set of motivated cycles naturally forms a graded \mathbb{Q} -algebra. The category of motives over K , denoted Mot_K , consists of objects (X, p, m) , where X is a smooth projective variety over K , p is an idempotent motivated cycle on $X \times X$, and m is an integer. A morphism $(X, p, m) \rightarrow (Y, q, n)$ is a motivated cycle γ of degree $n - m$ on $Y \times X$ such that $q\gamma p = \gamma$. We denote with $H_{\text{mot}}(X)$ the object $(X, \Delta, 0)$, where Δ is the class of the diagonal in $X \times X$. The Künneth projectors π_i are motivated cycles, and we denote with $H_{\text{mot}}^i(X)$ the object $(X, \pi_i, 0)$. Observe that $H_{\text{mot}}(X) = \bigoplus_i H_{\text{mot}}^i(X)$. This gives contravariant functors $H_{\text{mot}}(-)$ and $H_{\text{mot}}^i(-)$ from the category of smooth projective varieties over K to Mot_K .

THEOREM 5.4. The category Mot_K is Tannakian over \mathbb{Q} , semisimple, graded, and polarised. Every classical cohomology theory of smooth projective varieties over K factors via Mot_K .

Proof. See théorème 0.4 of [And96]. □

Definition 5.5. Let K be a subfield of \mathbb{C} . An *abelian motive* over K is an object of the Tannakian subcategory of Mot_K generated by objects of the form $H_{\text{mot}}(X)$ where X is either an Abelian variety or $X = \text{Spec}(L)$ for some finite extension L/K , with $L \subset \mathbb{C}$.

We denote the category of abelian motives over K with AbMot_K .

Finally we need the following theorem

THEOREM 5.6. The Hodge realization functor $H(-)$ restricted to the subcategory of abelian motives is a full functor.

Proof. See théorème 0.6.2 of [And96]. □

By Theorem 5.4, the singular cohomology and ℓ -adic cohomology functors factor via Mot_K . This means that if M is a motive, then we can attach to it a Hodge structure $H(M)$ and an ℓ -adic Galois representation $H_\ell(M)$. The Artin comparison isomorphism between singular cohomology and ℓ -adic cohomology extends to an isomorphism of vector spaces $H_\ell(M) \cong H(M) \otimes \mathbb{Q}_\ell$ that is natural in the motive M .

We can state the motivated Mumford–Tate conjecture following [Moo17b, Section 3.2]. We shall write $G_{\text{MT}}(M)$ for the Mumford–Tate group $G_{\text{MT}}(H(M))$. Similarly, we write $G_\ell(M)$ (resp. $G_\ell^\circ(M)$) for $G_\ell(H_\ell(M))$ (resp. $(G_\ell^\circ(H_\ell(M)))$) for the Tate group. The Mumford–Tate conjecture extends to motives: for the motive M it asserts that the comparison isomorphism $H_\ell(M) \cong H(M) \otimes \mathbb{Q}_\ell$ induces an isomorphism

$$G_\ell^\circ(M) \cong G_{\text{MT}}(M) \otimes \mathbb{Q}_\ell.$$

The discussion we have given here we enable us to prove the following.

PROPOSITION 5.7. Let S be a surface of general type as above if there exists a subgroup H of the automorphisms group G , and there exist a K3 surface X birational to S/H and ι as in (H) is *motivated* (in the sense of André) then the Tate and Mumford–Tate conjectures hold for S . That is the Tate and Mumford–Tate conjectures hold for those S that are deformation equivalent to such a surface S .

Proof. The proof of this theorem is contained in [CoPe20, Section 5]. We illustrate only the demonstration strategy in the realm of motives.

The main idea in [CoPe20] is that for surfaces S with $p_g = 2$ it is sometimes possible to decompose the weight 2 Hodge structure into two Hodge substructures of K3 type and see that these Hodge substructures are indeed the Hodge structures of either Abelian surfaces or K3 surfaces which are (birational) quotients of S . This geometric construction makes possible to consider the theory of motivated cycles introduced by André, and to decompose the motive of S into two abelian motives of K3 type. For these motives the Mumford–Tate conjecture is known. This, together with the main results of [Com16] and [Com19] allows to prove the Mumford–Tate conjecture for S . In [CoPe20] it was used the fact that the surfaces studied are of maximal Albanese dimension hence there is naturally an Abelian surface as a quotient surface. □

Let us summarize here, in the form of a table, the classification of the minimal surfaces of general type with $p_g = q = 2$ and of maximal Albanese dimension. Moreover, for each family we will indicate whether the Mumford–Tate conjecture has been proved or not.

In Table 1 have indicated, where possible, the number of families (#) and the dimensions of the irreducible component containing them (dim). Moreover, we point out if some members of the family are product-quotient surfaces (PQ) or mixed surfaces (MS), in particular for some PP2 surfaces the proof is given by the second author in [Pig20], while for the remaining ones is unknown. In the last column, we give references to more detailed descriptions of the class. Finally, we put a checkmark in the column MTC if the strategy given in [CoPe20] is enough to prove the Tate and Mumford–Tate conjectures for a class. As one can see, up to now, it is the only way to prove Mumford–Tate conjecture for these surfaces.

N ^o	K_S^2	$\deg(\alpha)$	#	dim	Name	MTC	PQ/MS	Reference
1	8	2	2	0^2			No	[PRR20]
2	8	{2,4,6}	4	$3^3, 4$	SIP	✓	Yes	[Pen11]
3	7	3	1	3	PP7	✓	Yes	[PiPo17, CaFr18]
4	7	2	3	3			?	[Rit18, PePi20]
5	6	4	1	4	PP4	✓	Yes	[PePo14]
6	6	3	?	3	AC3		?	[AlCa22, CaSe22]
7	6	2	1	3	PP2		?	[PePo13b]
8	6	2	2	4^2	PP2	✓	Yes	[PePo13b] [Pig20]
9	5	3	1	4	CHPP	✓	Yes	[ChHa06, PePo13a]
10	4	2	1	4	Catanese	✓	Yes	[Pen11, CiML02]

Table 1: State of the art of the classification of minimal complex algebraic surfaces with invariants $p_g = q = 2$ and with maximal Albanese dimension.

Finally, we explain here the meaning of the half-line between the surfaces with $K_S^2 \geq 7$ and $K_S^2 \leq 6$. This line is due to the classification theorem of [DJZ23], who recently proved that the classification of surfaces with $p_g = q = 2$ and $K_S^2 \leq 6$ is complete.

References

- [AlCa22] M. Alessandro, F. Catanese *On the components of the Main Stream of the moduli space of surfaces of general type with $p_g = q = 2$* , preprint available at arXiv:2212.14872
- [And96] Y. André *Pour une théorie inconditionnelle des motifs*. In: Institut des Hautes Études Scientifiques. Publications Mathématiques **83** (1996), pp. 5 – 49.
- [Bar87] W. Barth: *Abelian surfaces with (1,2)–Polarization*. Algebraic Geometry, Sendai, 1985, 41–84, Adv. Stud. Pure Math. **10**, North-Holland, Amsterdam, 1987.
- [BHPV03] W. Barth, K. Hulek, C.A.M. Peters, A. Van de Ven, *Compact Complex Surfaces*. Grundlehren der Mathematischen Wissenschaften, Vol **4**, Second enlarged edition, Springer-Verlag, Berlin, 2003.
- [CaFr18] N. Cancian, D. Frapporti, *On semi-isogenous mixed surfaces*, Math. Nachr. **291**, 264–283 (2018).
- [CaSe22] F. Catanese, E. Sernesi, *The Hesse pencil and polarizations of type (1,3) on Abelian surfaces*, preprint available at arXiv:2212.14877.
- [ChHa06] J. A. Chen, C. D. Hacon, *A surface of general type with $p_g = q = 2$ and $K_X^2 = 5$* , Pacif. J. Math. **223** (2006), 219–228.
- [CiML02] C. Ciliberto, M. Mendes Lopes, *On surfaces with $p_g = q = 2$ and non-birational bicanonical maps*, Advances in Geometry **2.3** (2002), 281–300.
- [Com16] J. Commelin, *The Mumford–Tate conjecture for the product of an abelian surface and a K3 surface*, Documenta Mathematica **21** (2016), 1691–1713.
- [Com19] J. Commelin, *The Mumford–Tate conjecture for products of abelian varieties*, Algebr. Geom. **6** (2019), 650–677.
- [CoPe20] J. Commelin, M. Penegini, *On the cohomology of surfaces with $p_g = q = 2$ and maximal Albanese dimension*, Transactions of the American Mathematical Society, 2020, **373**, 1749–1773.

- [DeMu91] C. Deninger and J. P. Murre, *Motivic decomposition of abelian schemes and the Fourier transform*, in: J. Reine und Angew. Math. **422** (1991), 201–219.
- [DJZ23] J. Du, Z. Jiang, G. Zhang, *Cohomological rank functions and surfaces of general type with $pg=q=2$* , preprint available at ArXiv:2309.05097
- [Lat19] R. Laterveer, *Algebraic cycles and triple K3 burgers* Ark. Mat. **57**(1) (2019), 157–189.
- [Man68] J. I. Manin, *Correspondences, motifs and monoidal transformations*. In: Mat. Sb. (N.S.) **77** (1968), 475–507.
- [Moo17a] B. Moonen, *On the Tate and Mumford–Tate conjectures in codimension 1 for varieties with $h^{2,0} = 1$* . Duke Mathematical Journal **166** (2017), 739–799.
- [Moo17b] B. Moonen, *Families of Motives and the Mumford–Tate Conjecture*, Milan J. Math. **85**, 257–307 (2017).
- [Mor87] D. R. Morrison. *Isogenies between algebraic surfaces with geometric genus one*. In: Tokyo Journal of Mathematics **10.1** (1987), pp. 179–187.
- [Mur90] J. P. Murre, *On the motive of an algebraic surface*, in: J. Reine und Angew. Math. **409** (1990), 190–204.
- [Pen11] M. Penegini, *The classification of isotrivially fibred surfaces with $p_g = q = 2$* . Collect. Math. **62**, 239–274 (2011).
- [PePi20] M. Penegini, R. Pignatelli, *A note on a family of surfaces with $p_g = q = 2$ and $K^2 = 7$* , Boll. Unione Mat. Ital. **15**, 305–331 (2022).
- [PePo13a] M. Penegini, F. Polizzi, *On surfaces with $p_g = q = 2$, $K^2 = 5$ and Albanese map of degree 3*, Osaka J. Math. **50**, 643–686 (2013).
- [PePo13b] M. Penegini, F. Polizzi, *On surfaces with $p_g = q = 2$, $K^2 = 6$ and Albanese map of degree 2*, Canad. J. Math. **65** (2013), 195–221
- [PePo14] M. Penegini, F. Polizzi, *A new family of surfaces with $p_g = q = 2$ and $K^2 = 6$ whose Albanese map has degree 4*, J. London Math. Soc. **90**, 741–762 (2014).
- [Pig20] R. Pignatelli, *Quotients of the square of a curve by a mixed action, further quotients and Albanese morphisms*. Revista Matematica Complutense **33** (2020), 911–931.
- [PiPo17] R. Pignatelli, F. Polizzi *A family of surfaces with $p_g = q = 2$, $K^2 = 7$ and Albanese map of degree 3* Mathematische Nachrichten, 2017, **290**, pp. 2684–2695
- [PRR20] F. Polizzi, C. Rito, X. Roulleau, *A pair of rigid surfaces with $p_g = q = 2$ and $K^2 = 8$ whose universal cover is not the bidisk*, Int. Math. Res. Not. IMRN 2020, **11**, 3453–3493 (2020).
- [Rit18] C. Rito *New surfaces with $K^2 = 7$ and $p_g = q \leq 2$* , Asian J. Math. **22** (2018), no. 6, 1117–1126.
- [Sil08] J.H. Silverman, *The Arithmetic of Elliptic Curves*, 2nd Edition, Springer GTM vol **106**, (2008).

Matteo Penegini, Università degli Studi di Genova, DIMA Dipartimento di Matematica, I-16146 Genova, Italy
e-mail penegini@dimma.unige.it

Roberto Pignatelli Università degli Studi di Trento, Dipartimento di Matematica, I-38123 Trento, Italy
e-mail Roberto.Pignatelli@unitn.it

AMS Subject Classification: 14J29, 14J10, 14B12

Lavoro pervenuto in redazione il MM.GG.AAAA.